

Preventing blood clots in pregnancy and after giving birth

Maternity Department
Patient Information Leaflet



Accredited as
Baby Friendly by

unicef 

Introduction

This leaflet is for pregnant women. It gives information about blood clots in pregnancy, what causes them and what you can do to help prevent them.

What is a venous thromboembolism (VTE)?

Venous thromboembolism (VTE) is a term for both deep vein thrombosis (DVT) and pulmonary embolism (PE).

Deep vein thrombosis (DVT)

A DVT is a blood clot in the deep veins of the leg or pelvis. It can block off or reduce the flow of blood in a vein. It can cause a condition called postthrombotic syndrome which can result in chronic pain and swelling of the leg.

Pulmonary embolism (PE)

A PE is when all or part of the blood clot breaks off, travels through the body and blocks the blood vessels to the lungs. It is potentially life threatening.

What are the signs and symptoms of VTE?

- Swelling, usually in one leg. If the thrombosis is in the thigh veins as is most common during pregnancy and following child birth, the whole leg may be swollen
- Tenderness of the leg muscles, often the calf muscles
- Reddish or bluish skin discoloration
- A leg that is warm to touch
- Sudden shortness of breath
- Chest pain that is sharp or stabbing and that may get worse when breathing in
- Rapid heart rate
- Unexplained cough, sometimes with bloody mucus
- Collapse (in severe cases)

If you think you may have DVT or PE, you should seek medical advice immediately either from your GP or nearest A&E (Emergency Department).

What causes VTE in pregnancy and after giving birth?

There are many changes in your body during pregnancy. One of these is that you are at increased risk of developing blood clots. Your blood becomes stickier and the blood flow slows down in the veins in your legs and pelvis which causes this increased risk. It can take around six weeks after you give birth for your blood to return to normal.

There are other factors that may increase your risk of a blood clot. These include:

- Having VTE previously
- A family history of VTE
- Assisted conception (IVF)
- Multiple pregnancies (twins or more)
- Age 35 years or older
- Weight (BMI above 30)
- If you already have three or more children
- Being sick a lot during your pregnancy
- Certain illnesses such as cancer, heart failure, lung disease, inflammatory diseases, infections such as pneumonia
- Smoking
- Not moving around much; for example, if you are unwell in bed or you have a leg in a plaster cast
- Long distance travel for more than four hours by car, train, coach or aeroplane, if you do not move around
- A long labour
- Delivery by caesarean section

Risk assessment for blood clots

Who will be risk assessed?

All pregnant woman will be assessed for potential risk of a blood clot developing.

When will I be assessed?

The assessment will be carried out by a midwife or a doctor at your booking appointment.

If you are identified as high risk for developing blood clots, we will refer you to a specialist clinic, for treatment to help reduce this risk.

This risk assessment will be reviewed throughout your pregnancy and immediately after the birth of your baby.

How will I be assessed?

We will assess how you are during pregnancy and ask about your medical and family history.

What treatment is there to help prevent VTE?

If you are considered to be at high risk of developing a blood clot, you will be offered treatment to help prevent one from forming.

You may be offered medication to thin your blood. This is called heparin and is given every day by injection.

In certain circumstances, we may recommend anti-embolism stockings. These are made from elastic and can be either thigh or knee length. They reduce the risk of blood clots by gently compressing your legs. This increases the blood flow and prevents your leg veins from expanding, which stops blood pooling in your legs and forming a clot.

If you are given them, you should wear the anti-embolism stockings while you are in hospital to give birth and until you can move around normally.

Do these treatments have side effects?

There are minor side effects from heparin. You may find you have bruising around the injection site. If you need to have heparin, all potential side effects will be discussed with you before the treatment begins.

Will the treatment affect my baby?

Heparin does not cross the placenta so it is safe for your developing baby and it is safe if you are breastfeeding.

What else can I do to help prevent VTE?

- Move around and try not to sit still for too long.
- On long journeys, try to move your legs and where possible, have a break and walk around every one to two hours.
- Drink plenty of fluids and do not become dehydrated.
- If you have a caesarean section, getting up and about as soon as possible after the caesarean can reduce the risk of VTE.
- It is essential that if you start having heparin treatment, you complete the full course.

What happens when I go home after giving birth?

Tell your doctor or midwife if you experience any of the following on your feet or legs:

- Strange marks on your skin
- Blistering
- Pain or discomfort
- Discoloration of skin

If you need to have heparin injections at home, your midwife will show you how to do this. The midwife will also give you a sharps bin to dispose of the used needles safely. If you need additional help, please ask your midwife.

You can return your used sharps bin to the ward that gave it to you. In addition, some GP surgeries and pharmacies may accept it.

If you need anti-embolism stockings, we measure and fit you with these. We will show you how to wear the stockings correctly. You should remove your stockings every day so you can wash your legs and check your skin for any changes.

Please refer to the manufacturer's instructions for washing the stockings and for other handling advice.

Can I find out more?

You can find out more from the following weblink:

NHS Choices

<http://www.nhs.uk/conditions/pregnancy-and-baby/pages/dvt-blood-clot-pregnant.aspx>

If you have any questions, or if there is anything you do not understand about this leaflet, please contact:

Maternity triage on 01384 456111 ext. 3053

This leaflet can be downloaded or printed from:

<http://dudleygroup.nhs.uk/services-and-wards/maternity/>

This leaflet can be made available in large print, audio version and in other languages, please call 0800 073 0510.

للحصول على هذه النشرة بحجم أكبر، وعلى شكل إصدار صوتي و بلغات أخرى، الرجاء الاتصال بالرقم 08000730510.

此宣传单可提供大字版本、音频版本和其它语言版本，请拨打电话：0800 073 0510。

Ulotka dostępna jest również w dużym druku, wersji audio lub w innym języku. W tym celu zadzwoń pod numer 0800 073 0510.

ਇਹ ਪਰਚਾ ਵੱਡੇ ਅੱਖਰਾਂ, ਬੋਲ ਕੇ ਰੀਕਾਰਡ ਕੀਤਾ ਹੋਇਆ ਅਤੇ ਦੂਸਰੀਆਂ ਭਾਸ਼ਾਵਾਂ ਵਿਚ ਵੀ ਪ੍ਰਾਪਤ ਹੋ ਸਕਦਾ ਹੈ, 0800 073 0510 ਤੇ ਫੋਨ ਕਰੋ ਜੀ।

Aceasta brosură poate fi pusă la dispoziție tipărită cu caractere mari, versiune audio sau în alte limbi, pentru acest lucru va rugăm sunați la 0800 073 0510.

یہ کتابچہ آپ کو بڑے حروف کی لکھائی، سمعی صورت اور دیگر زبانوں میں مہیا کیا جا سکتا ہے۔ برائے مہربانی فون نمبر 08000730510 پر رابطہ کریں۔